

1. Sato E, Mori T, Mishima E, Suzuki A, Sugawara S, Kurasawa N, Saigusa D, Miura D, Morikawa-Ichinose T, Saito R, Oba-Yabana I, Oe Y, Kisu K, Naganuma E, Koizumi K, Mokudai T, Niwano Y, Kudo T, Suzuki C, Takahashi N, Sato H, Abe T, Niwa T, Ito S. Metabolic alterations by indoxyl sulfate in skeletal muscle induce uremic sarcopenia in chronic kidney disease. *Sci Rep.* 2016;6:36618. doi: 10.1038/srep36618.
2. Torigoe A, Sato E, Mori T, Ieiri N, Takahashi C, Ishida Y, Hotta O, Ito S. Comparisons of amino acids, body constituents and antioxidative response between long-time HD and normal HD. *Hemodial Int.* 2016;20 Suppl 1:S17-S24. doi: 10.1111/hdi.12462.
3. Sato E, Mori T, Satoh M, Fujiwara M, Nakamichi Y, Oba I, Ogawa S, Kinouchi Y, Sato H, Ito S, Hida W. Urinary angiotensinogen excretion is associated with blood pressure in obese young adults. *Clin Exp Hypertens.* 2016;38(2):203-8. doi: 10.3109/10641963.2015.1081219.
4. Shimada S, Mori T, Koizumi K, Sato S, Oba-Yabana I, Ohsaki Y, Sato E, Naganuma E, Kurasawa N, Tsuchikawa M, Ito S. Efficacy and Biocompatibility of Neutral Icodextrin Peritoneal Dialysis Fluid. *Adv Perit Dial.* 2016;32:46-50.
5. Mori T, Kurasawa N, Ohsaki Y, Koizumi K, Sato S, Oba-Yabana I, Shimada S, Sato E, Naganuma E, Tsuchikawa M, Ito S. Role of Chronic Use of Tolvaptan in Patients with Heart Failure Undergoing Peritoneal Dialysis. *Adv Perit Dial.* 2016;32:39-45.
6. Tojo A, Kinugasa S, Fujita T, Wilcox CS. A local renal renin-angiotensin system activation via renal uptake of prorenin and angiotensinogen in diabetic rats. *Diabetes Metab Syndr Obes.* 2016;9:1-10. doi: 10.2147/DMSO.S91245.
7. Tojo A, Asaba K, Kinugasa S, Ikeda Y, Shintani Y, Fukayama M, Nangaku M. The reduced expression of proximal tubular transporters in acquired Fanconi syndrome with κ light chain deposition. *Med Mol Morphol.* 2016;49(1):48-52. doi: 10.1007/s00795-015-0114-3.